

SUBLINGUAL IMMUNOTHERAPY TREATMENT SET

Your sublingual immunotherapy treatment (SLIT) set consists of 1 x 12 ml Starter Dose vial (1 month supply) and 1 x 32 ml or 2 x 32 ml Maintenance Dose vial(s) (4 months supply each). You will use the Starter Dose vial for the first month or what is called the “build up phase” of oral allergy treatment, followed by the Maintenance Dose vial(s) for approximately four to eight months.

A 2-vial treatment set lasts a total of 5 months.

A 3-vial treatment set lasts a total of 9 months.

Suggested Dosage Schedule

Using the Starter Dose Vial (small yellow label vial) administer **3 sprays** into side of mouth or under the tongue, 7 days/week for 4 weeks.

NOTE START DATE HERE: _____

After for weeks, begin dosing with the Maintenance Dose (Large green label vial), administering **3 sprays** into side of mouth or under the tongue, 6 days/week. (Each vial will last approximately 4 months.) Use each vial until empty.

	DAY	DOSE
<u>Build-up Phase</u> Use Starter Vial (yellow label)	1 to 28	3 "squirts" per day or approximately 0.3 ml per day
	DAY	DOSE*
<u>Maintenance Phase</u> Use Maintenance Vial (green label) Note: Use Maintenance Vials one at a time until each is empty.	From day 29 forward	3-8- "squirts" per day six days per week

*The maintenance dose is a dose that stimulates positive changes in the animal’s immune system but does not cause clinical signs or symptoms to recur. The maintenance dose may need adjustment depending on the animal’s response.

IMPORTANT – TO AVOID INTERRUPTING YOUR PET’S TREATMENT, CONTACT YOUR VETERINARIAN TO REORDER.

STORAGE

Refrigeration at 2°C to 8°C and no higher than 46°F (8°C) recommended. DO NOT FREEZE.

SPECIAL PRECAUTIONS:

- 1. DO NOT INJECT THIS MATERIAL INTO THE ANIMAL.**
2. To aid absorption of allergenic extract, the animal should be kept from food or drink for 30 minutes before and after receiving a dose.
3. Therapy should be started with the Starter Vial, which is the weaker solution.
4. Oral allergy treatment (SLIT) is given by dispensing the allergenic extract under the animal’s tongue.
5. Animal should be observed for untoward reactions for 30 minutes after receiving a dose.
6. *In the unlikely event that a human might receive a dose of this product, immediately consult a physician experienced in allergic reaction. If a severe allergic reaction occurs (i.e.: shock, anaphylaxis, dyspnea) administer oral antihistamines or intramuscular epinephrine and go to an emergency center.*